



Specific Changes in Young Soccer Player's Fitness After Traditional Bilateral vs. Unilateral Combined Strength and Plyometric Training

Rodrigo Ramirez-Campillo^{1,2}, Javier Sanchez-Sanchez², Oliver Gonzalo-Skok^{2,3}, Alejandro Rodríguez-Fernández^{2,4}, Manuel Carretero² and Fabio Y. Nakamura^{2,5,6*}

OPEN ACCESS

Edited by:

David George Behm,
Memorial University of Newfoundland,
Canada

Reviewed by:

Pantelis Theodoros Nikolaidis,
Hellenic Army Academy, Greece
Anis Chaouachi,
National Center of Medicine and
Science in Sports, Tunisia
Thomas Muehlbauer,
University of Duisburg-Essen,
Germany
Jonathan Peter Farthing,
University of Saskatchewan, Canada

*Correspondence:

Fabio Y. Nakamura
fabioy_nakamura@yahoo.com.br

Specialty section:

This article was submitted to
Exercise Physiology,
a section of the journal
Frontiers in Physiology

Received: 02 October 2017

Accepted: 08 March 2018

Published: 22 March 2018

Citation:

Ramirez-Campillo R,
Sanchez-Sanchez J, Gonzalo-Skok O,
Rodríguez-Fernández A, Carretero M
and Nakamura FY (2018) Specific
Changes in Young Soccer Player's
Fitness After Traditional Bilateral vs.
Unilateral Combined Strength and
Plyometric Training.
Front. Physiol. 9:265.
doi: 10.3389/fphys.2018.00265

¹ Department of Physical Activity Sciences, Research Nucleus in Health, Physical Activity and Sport, Universidad de Los Lagos, Osorno, Chile, ² Research Group Planning and Assessment of Training and Athletic Performance, Pontifical University of Salamanca, Salamanca, Spain, ³ Faculty of Health Sciences, University of San Jorge, Zaragoza, Spain, ⁴ Facultad de Ciencias de la Salud, Universidad Isabel I, Burgos, Spain, ⁵ The College of Healthcare Sciences, James Cook University, Townsville, QLD, Australia, ⁶ Department of Medicine and Aging Sciences, "G. d'Annunzio" University of Chieti-Pescara, Chieti, Italy

The aim of this study was to compare changes in young soccer player's fitness after traditional bilateral vs. unilateral combined plyometric and strength training. Male athletes were randomly divided in two groups; both received the same training, including strength training for knee extensors and flexors, in addition to horizontal plyometric training drills. The only difference between groups was the mode of drills technique: unilateral (UG; $n = 9$; age, 17.3 ± 1.1 years) vs. bilateral (TG; $n = 9$; age, 17.6 ± 0.5 years). One repetition maximum bilateral strength of knee muscle extensors (1RM_KE) and flexors (1RM_KF), change of direction ability (COD), horizontal and vertical jump ability with one (unilateral) and two (bilateral) legs, and limb symmetry index were measured before and after an 8-week in-season intervention period. Some regular soccer drills were replaced by combination of plyometric and strength training drills. Magnitude-based inference statistics were used for between-group and within-group comparisons. Beneficial effects ($p < 0.05$) in 1RM_KE, COD, and several test of jumping performance were found in both groups in comparison to pre-test values. The limb symmetry index was not affected in either group. The beneficial changes in 1RM_KE (8.1%; $p = 0.074$) and 1RM_KF (6.7%; $p = 0.004$), COD (3.1%; $p = 0.149$), and bilateral jump performance (from 2.7% [$p = 0.535$] to 10.5% [$p = 0.002$]) were possible to most likely beneficial in the TG than in the UG. However, unilateral jump performance measures achieved likely to most likely beneficial changes in the UG compared to the TG (from 4.5% [$p = 0.090$] to 8.6% [$p = 0.018$]). The improvements in jumping ability were specific to the type of jump performed, with greater improvements in unilateral jump performance in the UG and bilateral jump performance in the TG. Therefore, bilateral strength and plyometric training should be complemented with unilateral drills, in order to maximize adaptations.

Keywords: team-sports, football, strength, change of direction ability, young athletes